

## DAFTAR PUSTAKA

- A. Ceriello, “Oxidative stress and glycemic regulation,” *Metabolisme Clinical and Experimental*, vol. 49, no. 2, pp. 27–29, 2000.
- Adisakwattana, S.; Chanathong, B. Alpha-glucosidase inhibitory activity and lipid-lowering mechanisms of *Moringa oleifera* leaf extract. *Eur. Rev. Med. Pharmacol. Sci.* **2011**, 15, 803–808.
- Adisakwattana, S.; Chanathong, B. Alpha-glucosidase inhibitory activity and lipid-lowering mechanisms of *Moringa oleifera* leaf extract. *Eur. Rev. Med. Pharmacol. Sci.* 2011, 15, 803–808.
- Alam MN, Bristi NJ, Rafiquzzaman M (2013) Review on in vivo and in vitro methods evaluation of antioxidant activity. *Saudi Pharm J* 21: 143-152.
- Alkhaldy, H.; Moore, W.; Zhang, Y.; McMillan, R.; Wang, A.; Ali, M.; Suh, K.- S.; Zhen, W.; Cheng, Z.; Jia, Z. Small molecule kaempferol promotes insulin sensitivity and preserved pancreatic b-cell mass in middle-aged obese diabetic mice. *J. Diabetes Res.* **2015**, 2015, 532984.
- American Diabetes Association, 2014. *Diagnosis and classification of diabetes mellitus. Diabetes Care* 37 (Suppl. 1), S81–S90.
- Anwar, F.; Latif, S.; Ashraf, M.; Gilani, A.H. *Moringa oleifera*: A food plant with multiple medicinal uses. *Phytother. Res.* **2007**, 21, 17–25.
- Annunziata F, Pinna C, Dallavalle S, Tamborini L, Pinto A. (2020). An overview of coumarin as a versatile and readily accessible scaffold with broad-ranging biological activities. *Int J Mol Sci*, 21, 1-83.
- Arjadi, F., & Susatyo, P. (2010). Islet of langerhans regeneration in diabetic white rats (*Rattus norvegicus*) after giving decocted pulp of mahkota dewa. (*Phaleria macrocarp* (scheff.) Boerl), 2(2), 117-126.
- Badan Pengawas Obat dan Makanan Republik Indonesia. (2006). *Monografi ekstrak tumbuhan obat Indonesia*. (Volume 2). Jakarta: Badan Pengawas Obat dan Makanan Republik Indonesia.

- Bajaj S. and Khan A. Antioxidant and diabetes. Indian Journal of Endocrinology and Metabolism. 2012;16(2): 267-271a.
- Benzie, I.F.F., Strain, J.J., 1996. The ferric reducing ability of plasma (FRAP) as a measure of 'antioxidant power': The FRAP assay. *Anal. Biochem.* 239, 70–76.
- Bour, S., Visentin, V., Prevot, D., Daviaud, D., Saulnier-Blache, J. S., Guigne, C., Valet, P., and Carpene, C. (2005). Effects of oral administration of benzylamine on glucose tolerance and lipid metabolism in rats. *J. Physiol. Biochem.* 61, 371–379.
- Brunner & Suddarth, Buku Ajar Keperawatan Medikal Bedah Edisi 8 volume 2. Jakarta. EGC.2013
- C. Rosak and G. Mertes, "Effects of acarbose on proinsulin and insulin secretion and their potential significance for the intermediary metabolism and cardiovascular system," *Current Diabetes Reviews*, vol. 5, no. 3, pp. 157– 164, 2009.
- Ceriello A and Testa R. Antioxidant Anti-Inflammatory Treatment in Type 2 Diabetes. *Diabetes Care.* 2009;32(2): 5232-5236.
- Chakravarti, R. N. (1955). Chemical identity of moringine. *Bull. Calcutta Sch.Trop. Med.* 3, 162–163.
- Chang, C. W., S.C ., Hwang SS., Choi, B. K., Ahn, H. J., Lee, M. Y., Park, S. H., & Kim, S. K. Antioxidant Activity and Free Radical Scavenging Capacity Between Korean medicinal plants dan Flavonoids by Assay-guide comparison. *Plant Science*, 163: 1161-1168.
- Chen A, Chen Y. A review of the dietary flavonoid, kaempferol on human health and cancer chemoprevention. *Food Chem.* 2013;138(4):2099–107.
- Choi CW, Kim SC, Hwang SS, Choi BK, Ahn HJ, Lee MY, Park SH, Kim SK. Antioxidant activity and free radical scavenging capacity between Korean medicinal plants and flavonoids by assay-guided comparison. *Plant Sci* 2002;163:1161-8.
- Dalimarta S. *Atlas tumbuhan obat Indonesia Jilid I.* Jakarta: Trubus Agriwidya; 2014.
- DD Chakraborty, V Ravi, P Chakraborty, Phytochemical evaluation

- and TLC protocol of various extracts of *Bombax ceiba* Linn, *International Journal of Pharmaceutical Sciences and Research*, 1, 8, (2010) 66-73.
- Deng, S., M. Vatamanjuk, X. Huang, N. Doliba, M.M. Lian, A. Frank, E. Velidededeoglu And F. Markmamm. 2004. Structural and functional abnormalities in the islets isolated from type 2 diabetic subjects. *Diabetes* 53: 624-632.
- Departemen Kesehatan Republik Indonesia. 1989. *Materia Medika Indonesia* Edisi V. Direktorat Jenderal Pengawasan Obat dan Makanan, Jakarta.
- Departemen Kesehatan Republik Indonesia. 2008. *Farmakope Herbal Indonesia*. Jakarta: Departemen Kesehatan Republik Indonesia.
- Departemen Kesehatan Republik Indonesia. 2017. *Farmakope Herbal Indonesia*. Edisi II. Jakarta: Departemen Kesehatan Republik Indonesia.
- Diani, A.R., G. Sawada, B. Wyse, F.T. Murray And M. Khan. 2004. Pioglitazone preserves pancreatic islet structure and insulin secretory function in three murine models of type 2 diabetes. *Am. J. Physiol. Endocrinol. Metab.* 286: 116-122.
- Duo J, Ying G, Wang G, Zhang L. Quercetin inhibits human breast cancer cell proliferation and induces apoptosis via Bcl-2 and Bax regulation. *Molecular Medicine Reports*. 2012; 5(6): 1453-6.
- E. Nicolle, F. Souard, P. Faure, and A. Boumendjel, “Flavonoids as promising lead compounds in type 2 diabetes mellitus: molecules of interest and structure-activity relationship,” *Current Medicinal Chemistry*, vol. 18, no. 17, pp. 2661–2672, 2011.
- Eid, H.M.; Nachar, A.; Thong, F.; Sweeney, G.; Haddad, P.S. The molecular basisof the antidiabetic action of quercetin in cultured skeletal muscle cells and hepatocytes. *Pharmacogn. Mag.* 2015, 11, 74–81. [PubMed].
- Fahey, J.W. 2005. *Moringa oleifera: A review of the Medical Evidence for Its Nutritional, Therapeutic and Prophylactic Properties*. Trees for Life Journal 2005, 1-5.

- Gargiulo .P, et al., *The metabolic syndrome in heart failure: insights to specific mechanisms*, Heart Fail. Rev. (2019).
- Gartner, Hiatt LP, Strum JL, et al. *Biologi Sel dan Histologi Edisi ke-6*. Jakarta: Binarupa Aksara Publisher. 2012.
- Gautam, M. K., Gangwar, M., Nath, G., Rao, CV & Goel, RK. (2012). Invitro antibacterial activity on human pathogens and total phenolic, flavonoid contents of *Murraya paniculata* Linn. Leaves. *Asian Pacific Journal of Tropical Biomedicine*, 2, (3), 1660-1663.
- Gautam, M.K., Gupta, A., Vijaykumar, M., Rao, CV & Goel, RK. (2012). Studies on the hypoglycemic effects of *Murraya Paniculata* Linn. Extract on alloxan-induced oxidative stress in diabetic and non-diabetic models.*Asian Pacific Journal of Tropical Disease*, 2, (1), 186-191.
- Ghorbani A, Rashidi R, Shafiee RN. Flavonoids for preserving pancreatic betacell survival and function: a mechanistic review. *Biomedicine & Pharmacotherapy*. 2019;111:947–57.
- Ghosh, S., Chopra, N. R., and Dutt, A. (1935). Chemical examination of bark of *Moringa pterygosperma*. Indian J. Med. Res. 22, 789.
- Guariguata, L., Whiting, D. R., Hambleton, I., Beagley, J., Linnenkamp, U., & Shaw, J. E. (2014). *Global estimates of diabetes prevalence for 2013 and projections for 2035*. *Diabetes Res Clin Pract*, 103(2), 137–149. <https://doi.org/10.1016/j.diabres.2013.11.002>.
- Gupta R, Mathur M, Bajaj VK, Katariya P, Yadav S, Kamal R, Gupta RS. Evaluation of antidiabetic and antioxidant activity of *Moringa oleifera* in experimental diabetes. *J Diabetes*. 2012;4:164–71.
- Gupta S, George M, Singhal M, et al. *Leaves extract of Murraya koenigii linn for anti-inflammatory and analgesic activity in animal models*. *J Adv Pharm Technol Res*; 2013.
- H.D. Yassa, A.F. Tohamy, *Extract of Moringa oleifera leaves ameliorates streptozotocin-induced Diabetes mellitus in adult rats*, *Acta Histochem*. 116 (5) (2014) 844–854.

Halliwell, B and Gutteridge, J.M.C., 2000, *Free Radical in Biology and Medicine*, Oxford University Press, New York.

Han, M.H.; Park, C.; Lee, D.S.; Hong, S.H.; Choi, I.W.; Kim, G.Y.; Choi, S.H.; Shim, J.H.; Chae, J.I.; Yoo, Y.H.; et al. Cytoprotective effects of esculetin against oxidative stress are associated with the upregulation of Nrf2-mediated NQO1 expression via the activation of the ERK pathway. *Int. J. Mol. Med.* **2017**, *39*, 380–386.

Herlina T, Tri H, Heru N, Ciptono (2018), Pengaruh Pemberian Ekstrak Biji Kedelai Putih (*Glycine Max, L.*) Terhadap Struktur Histologik Hati Dan Ginjal Tikus Putih (*Rattus Norvegicus, L.*), *128 Jurnal Prodi Biologi Vol 7No 2*.

Hevel JM, Marletta MA (1994) Nitric-oxide synthase assays. Methods Enzymol 233: 250-258.

Holt RIG, Cockram C, Flyvbjerg A & Goldstein BJ. *Textbook of Diabetes, 4th Edition*, Wiley-Blackwell, UK. 2010:160-207.

J. S. Johansen, A. K. Harris, D. J. Rychly, and A. Ergul, “Oxidative stress and the use of antioxidants in diabetes: linking basic science to clinical practice,” *Cardiovascular Diabetology*, vol. 4, article 5, 2005.

Jingtao Zou, Dayun Sui, Wenwen Fu, Yuangeng Li, Ping Yu, Xiaofeng Yu, Huali Xu, *Total flavonoids extracted from the leaves of Murraya paniculata (L.) Jack alleviate oxidative stress, inflammation and apoptosis in a rat model of diabetic cardiomyopathy*, Department of Pharmacology, School of Pharmaceutical Sciences, Jilin University, Changchun 130021, PR China, 2021.

Kandaswami, C and Middleton, E. 1997. Flavonoids as antioxidant, In F. Shahidi (Ed). *Natural Antioxidant Chemistry, Health Effects and Applications*. Champaign Illions : AOCS Press.

Katayoun S, Paydar M, Murphy B. *Pharmacological properties and chemical constituents of Murraya paniculata (L.) Jack*. Med Aromat Plants. 2014; 3(4).

Kementerian Kesehatan Republik Indonesia. 2011. *Formularium obat herbal asli Indonesia*. Jakarta: Kementerian Kesehatan Republik

Indonesia.

Kementerian Kesehatan Republik Indonesia, 2017. *Formularium ramuan obat tradisional indonesia*. Jakarta. Hal. 65.

Kementerian Kesehatan RI, (2020). *Infodatin: Tetap Produktif, cegah, dan atasi Diabetes Melitus*. Jakarta: Kementerian Kesehatan Republik Indonesia.

Kim M, Lee H, Lim B, Ha K, Kim S, So I, et al. Quercetin induces apoptosis by inhibiting MAPKs and TRPM7 channels in AGS cells. *Int J MolMed*. 2014;33(1):1657–63.

Kim, S., S. Jun-Seop, K. Hyun-Jung, K.C. Fisher, L. Mi-Ji and K.C Han-Wha. 2007. Streptozotocin-induced diabetes can be reversed by hepatic oval cell activation through hepatic transdifferentiation and pancreatic islet regeneration. *Lab. Investigation* 87: 702-712.

Kumbhare MR, Guleha V, Sivakumar T. Estimation of total phenolic content, cytotoxicity and in-vitro antioxidant activity of stem bark of *Moringa oleifera*. *Asian Pac J Trop Dis* 2012; 2(2): 144-150.

Lako, J., Trenerry, V. C., Wahlqvist, M., Wattanapenpaiboon, N., Sotheeswaran, S., and Premier, R. (2007). Phytochemical flavonols, carotenoids and the antioxidant properties of a wide selection of Fijian fruit, vegetables and other readily available foods. *Food Chem.* 101, 1727–1741.

Leone A, Spada A, Battezzati A, Schiraldi A, Aristil J, Bertoli S. Cultivation, genetic, ethnopharmacology, phytochemistry and pharmacology of *Moringa oleifera* leaves. An overview. *Int J Mol Sci.* 2015; 16(6): 12791– 835.

Liang, H.-J.; Suk, F.-M.; Wang, C.-K.; Hung, L.F.; Liu, D.-Z.; Che, N.-Q.; Chen, Y.-C.; Chang, C.-C.; Liang, Y.-C. Osthole, a potential antidiabetic agent, alleviates hyperglycemia in db/db mice. *Chem. Biol. Interact.* 2009, 181, 309–315.

Lobo V, Patil A, Phatak A, Chandra N. Free radicals, antioxidants and functional foods: impact on human health. *Pharmacogn Rev.* 2010;4:118.

- M. Mbikay, Therapeutic potential of *Moringa oleifera* leaves in chronichyperglycemia and dyslipidemia: a review, *Front. Pharmacol.* 3 (2012)1–12.
- Mescher LA. *Junquiera's Basic Histology Text and Atlas. 12<sup>th</sup> Edition.* New York: McGraw-Hill Companies, Inc. 2010.
- Mohan H. *Textbook of Pathology.* 5th ed. New Delhi: Jaypee Brothers Medical Publishers; 2010.
- N. F. Brás, N.M. F. Cerqueira, M. J. Ramos, and P. A. Fernandes, “Glycosidase inhibitors: a patent review (2008–2013),” *Expert Opinion on Therapeutic Patents*, vol. 24, no. 8, pp. 857–874, 2014.
- Narkhede, B.M., Ajmire, P.V & Wagh, A.E.(2012). Evaluation of antinociceptive and anti-inflammatory activity of ethanol extract of *Murraya Paniculata* leaves in experimentalrodents. *International Journal of Pharmacy and Pharmaceutical Science*, 4, (1), 247-250.
- Noro, T., Oda, Y., Miyase, T., Ueno, A., Fukushima, S., 1983. Inhibitors of xanthine oxidase from the flowers and buds of *Daphne genkwa*. *Chem. Pharm. Bull.* 31, 3984–3987.
- Nugroho A. B (2021), *Analisis Spektrum Tegangan Pada Alat Pendekripsi Kadar Gula Darah Menggunakan Near*, Jurnal Teknik Elektro dan Komputasi (ELKOM) Volume 3 Nomor 1 Maret 2021 || Hal. 1-13.
- Ogunwande, I.A.; Matsui, T.; Fujise, T.; Matsumoto, K. A-glucosidase inhibitory profile of nigerian medicinal plants in immobilized assay system. *Food Sci. Technol. Res.* **2007**, 13, 169–172.
- Och, M., Och, A., Cieśla, Ł., Kocki, J., & Bogucka-Kocka, A. (2013). *Screening various Juniperus species for the occurrence of umbelliferone by means of bivariate multiple development thin-layer chromatography.* JPC–Journal of Planar Chromatography–Modern TLC, 26(5), 421-426.
- Oyenihu AB, Ayeleso AO, Mukwevho E & Masola B. *Antioxidant strategies in the management of diabetic neuropathy.* BioMed Research International. 2015.

Ozougwu, JC., Obimba KC., Belonwu CD and Unakalamba CB. The Pathogenesis and Pathophysiology of type 1 and type 2 Diabetes Mellitus. *Journal of Physiology and Pathophysiology*. 2013;4(4): 46-57.

Puspitasari I. Uji aktivitas antihiperglikemik dan regenerasi sel pankreas fraksi-fraksi dari ekstrak biji mahoni (*swietenia macrophylla king*) pada tikus diabetes melitus yang diinduksi streptozotosin- nikotinamid. Tesis. Surakarta: Universitas Setia Budi, 2016.

Pokorny, J., Yanishlieva, N., and Gordon, M., 2001, *Antioxidant in Food; Practical Applications*, CRC Press, New York.

Rehman, F. U., Khan, M. F & Khan, I. (2014). Analgesic activity of carbazole alkaloid from *Murraya paniculata* Linn. (Rutaceae). *American Eurasian J. Agric & Environ.Sci*, 14, (3), 240-245.

Rivera, L., Moron, R., Sanchez, M., Zarzuelo, A., and Galisteo, M. (2008). Quercetin ameliorates metabolic syndrome and improves the inflammatory status in obese Zucker rats. *Obesity (Silver Spring)* 16, 2081–2087.

Rohman, A & Riyanto, S. (2005). Daya antioksidan eksrak etanol daun kemuning (*Murraya paniculata* (L.) Jack) secara *in vitro*. *Majalah Farmasi Indonesia*, 16, (3), 136-140.

Rohman, A. & Riyanto, S. 2005. Daya Antioksidan Ekstrak Daun Kemuning (*Murraya paniculata* (L) Jack) secara *in vitro*. *Majalah Farmasi Indonesia*, 16 (3): 136-140.

Rudianto, 2015, Studi Pembuatan Analisis Zat Gizi Pada Produk Biskuit Moringa Oleifera Dengan Subtitusi Tepung daun Kelor, Skripsi, Program Studi Ilmu Gizi FKM UH, Makassar.

S.M. Divi, R. Bellamkonda, S.K. Dasireddy, Evaluation of antidiabeticand antihyperlipidemic potential of aqueous extract of *Moringa oleiferain* fructose fed insulin resistant and STZ induced diabetic wistar rats: acomparative study, *Asian J. Pharm. Clin. Res.* 5 (2012) 67–72.

Sari, I. P., Pramono, S & Hali, A. A. (2004). Efek ekstrak etanolik daun kemuning (*Murraya paniculata* (L.) Jack) terhadap kolesterol

- darah tikus putih jantan diet tinggi lemak dan histopatologi hepar, jantung dan aorta, *Jurnal Lembaga Pendidikan Kepada Masyarakat UGM*, 1693, (1033), 44-46.
- Scheepers, A., Joost, H-G., Schurman, A. 2004. The Glucose Transporter Families SGLT and GLUT: Molecular Basis of Normal and Abberant Function, *Journal of Parenteral and Enteral Nutrition*, 28 (5): 364-371.
- Seeram, N.P., Henning, S.M., Lee, R., Niu, Y., Scheuller, H.S., Heber, D., 2006. Catechin and caffeine contents of green tea dietary supplements and correlation with antioxidant activity. *J. Agric. Food Chem.* 54, 1599–1603.
- Seidemann, J.2005. *World Spice Plants: Economic Usage, Botany, Taxonomy*. Berlin: Springer.
- Sharma, P., Batra, S., Kumar, A & Sharma, A. (2017). In vivo antianxiety and antidepressant activity of *Murraya Paniculata* leaf extracts. *Integrative Medicibne*, 15, (4), 320-325.
- Shin, E.; Choi, K.-M.; Yoo, H.-S.; Lee, C.-K.; Hwang, B.Y.; Lee, M.K. Inhibitory effects of coumarins from the stem barks of *Fraxinus rhynchophylla* on adipocyte differentiation in 3T3-L1 cells. *Biol. Pharm. Bull.* 2010, 33, 1610–1614.
- Sim, L., Willemsma, C., Mohan, S., Naim, H. Y., Pinto, M., & Rose D. R. 2010. Structural Basis for Substrate Selectivity in Human Maltase-Glucoamylase and Sucrase-Isomaltase N-terminal Domains. *The Journal of Biological Chemistry*, 285 (23): 17763-17770.
- Suarsana, I-N., B.P. Priosoeryanto, M. Bintang dan T. Wresdiyati. 2010. Profil glukosa darah dan ultrastruktur sel beta pankreas tikus yang diinduksi senyawa aloksan. *JITV* 15(2): 118-123.
- Sun, L., Peterson T.E., McCormick, M.L., Oberley, L.W., & Osborne, J.W. (1989). Improved superoxide dismutase assay for clinical use. *Clin Chem.*,35, 1265-1266.
- Surse, V.M.; Gupta, J.; Tikoo, K. Esculetin induced changes in Mmp13 and Bmp6 gene expression and histone H3 modifications attenuate development of glomerulosclerosis in diabetic rats. *J.*

- Mol. Endocrinol. 2011, 46, 245–254.
- Suyono S. *Kecenderungan Peningkatan Jumlah Penyandang Diabetes*. In: Sidartawan S, Pradana S, Imam S, editors. Penatalaksanaan Diabetes Mellitus. Jakarta: Badan Penerbit FK UI; 2011. p. 3-77.
- Szkudelski, T. 2001. The mechanism of alloxan and streptozotocin action in cells of the rat pancreas. *Physiol. Res.* 50: 536-546.
- Tadera, K.; Minami, Y.; Takamatsu, K.; Matsuoka, T. Inhibition of alpha- glucosidase and alpha-amylase by flavonoids. *J. Nutr. Sci. Vitaminol.* **2006**, 52, 149–153.
- Tallitsch RB, Guastaferri RS. *Histology: An Identification Manual*. Philadelphia: Mosby Elsevier. 2009. Chapter 12-13,16.
- Tandi J., Wulandari A., Asrifa. (2017). Efek Ekstrak Etanol Daun Gendola Merah (*Basella alba* L.) terhadap Kadar Kreatinin, Ureum dan Deskripsi Histologis Tubulus Ginjal Tikus Putih Jantan (*Rattus norvegicus*) Diabetes yang Diinduksi Streptozotocin. *Jurnal Farmasi Galenika (Galenika Journal of Pharmacy)*, 3(2), 93-102. doi:10.22487/j24428744.2017.v3.i2.8813.
- Tang, D.Q.; Wei, Y.Q.; Yin, X.X.; Lu, Q.; Hao, H.H.; Zhai, Y.P.; Wang, J.Y.; Ren, J. In vitro suppression of quercetin on hypertrophy and extracellular matrix accumulation in rat glomerular mesangial cells cultured by high glucose. *Fitoterapia* **2011**, 82, 920–926.
- Umar AN, Mann A, Ajiboso OSO. Ethnodiетetics of *Moringa oleifera* leaves amongst the ethnic groups in Bida, Niger State, Nigeria and its hypoglycaemic effects in rats. *Am Eurasian J Sustain Agric* 2011; 5(1): 107-14.
- V Steenis, CGGJ, 2008, *FLORA: untuk Sekolah di Indonesia*, Pradnya Paramita, Jakarta.
- Vongsak B, Sithisarn P, Gritsanapan W. HPLC quantitative analysis of three major antioxidative components of *Moringa oleifera* leaf extracts. *Planta Med* 2012; <http://dx.doi.org/10.1055/s-0032-1321175>.

- Whang WK, Park HS, Ham IH, et al. Natural compounds, fraxin and chemicals structurally related to fraxin protect cells from oxidative stress. *Exp Mol Med.* 2005;37(5):436-446. doi:10.1038/emm.2005.54.
- Widowati, Wahyu., Ratu Safitri, Rymond Rumumpuk, dan Marlinda Siahaan. 2005. *Penapisan Aktivitas Superoksida Dismutase pada Berbagai Tanaman.* JKM Vol. 5 (1): 33-47.
- Wood, J.L., 1970. In: Fishman, W.H. (Ed.), . In: Metabolic Conjugation and Metabolic Hydrolysis, vol. II. Academic Press, New York, pp. 61–299.
- Y. Zhenhua, Z. Wei, F. Fajin, Z. Yong, and K. Wenyi, “ $\alpha$ - Glucosidase inhibitors isolated from medicinal plants,” *Journal of Food Science and Human Wellness*, vol. 3, no. 3, pp. 136–174, 2014.
- Yang, G., Chen, J., Wen, Z., Ge, K., Zhu, L., Chen, X., 1984. The role of selenium in Keshan disease. *Adv. Nutr. Res.* 6, 203–220.
- Yongri J, Dayuan S, Xuwen L (2008) Applications of muraya jasmine or age leaf total flavones in preparation of medicament for curing diabetes. *Changchun Ruide Pharmaceutical Technology*.
- Youl, E.; Bardy, G.; Magous, R.; Cros, G.; Sejalon, F.; Virsolvay, A.; Richard, S.; Quignard, J.; Gross, R.; Petit, P.; et al. Quercetin potentiates insulin secretion and protects INS-1 pancreatic b-cells against oxidative damage via the ERK1/2. *Br. J. Pharmacol.* **2010**, 161, 799–814.
- Yu Lin, H. Kuo, Y.H. Lin, Y.L. dan Chiang, W., 2009, Antioxidative effect and active components from leaves of lotus (*Nelumbo nucifera*), *Journal of Agricultural and Food Chemistry* (57), 6623–6629.
- Zhang M, Hettiarachchy NS, Horax R, Kannan A, Apputhury Praisoody MD, Muhundan A, et al. Phytochemicals, antioxidant and antimicrobial activity of Hibiscus sabdariffa, Centella asiatica, Moringa oleifera and Murraya koenigii leaves. *J Med Plants Res* 2011; 5(30): 6672-80.
- Zhang Y, Jun L, Shepo S, Ke Z and Pengfei T. *Glycosides of Flavone Methyl Ethers from Murraya paniculata.* *Bioc. Syst. Eco* .2012;43(8): 10-13.