

DAFTAR PUSTAKA

- Liu R, Wang Y, Li J, dkk. Reduced T cell population contributes to increased severity COVID-19. *Clin Chim Acta*. 2020; 508: 110-114.
- Sit M, Aktas G, Erkol H, Yaman S, Keyif F, Savli H. Neutrophil about lymphocyte ratio is useful in malignant and benign thyroid nodules differentiation. *PR Kesehatan Sci J*. 2019; 38: 60-63.
- Jeong Y, Seong Jeon R, Kim HG, dkk. Role of platelet about lymphocyte ratio and neutrophil to lymphocyte ratio in ulcerative colitis. *Res usus*. 2020. <https://doi.org/10.5217/ir.2019.09156>.
- Aktas G, Sit M, Dikbas O, dkk. Increased neutrophil to lymphocyte ratio in Hashimoto's thyroiditis diagnosis. *Rev Assoc Med Bras*. 1992; 2017 (63): 1065-1068.
- Zeng Z, Ye W, Lin J, Ou Q. Red blood cell distribution width: potential laboratory parameters for monitoring inflammation in rheumatoid arthritis. *Clin Rheumatol*. 2018; 37: 161-167. Dagistan Y, Dagistan E, Gezici.
- AR, dkk. Can the width of red blood cell distribution and average platelet volume be a predictor of lumbar hernia disc? *Ideggyogy Sz*. 2016; 69: 411-414.
- Chen W, Zheng KI, Liu S, Yan Z, Xu C, Qiao Z. Plasma CRP levels are positively associated with severity COVID-19. *Ann Clin Microbiol Antimicrob*. 2020; 19 (1): 18.
- Liu M, Gao Y, Yuan Y, dkk. Efficacy and Safety of Integrated Traditional Chinese and Western Medicine for Coronavirus 2019 (COVID-19): tinjauan sistematis dan meta-analisis.
- Xu B, Fan CY, Wang AL, dkk. Suppressed T cell-mediated immunity in patients with COVID-19: a clinical retrospective study in Wuhan, China. *J infect*. 2020; 81: e51-e60.
- Zhou Y, Zhang Z, Tian J, Xiong S. Risk factors associated with the development of the disease in the cohort of patients infected with novel coronavirus 2019. *Ann Palliat Med*. 2020; 9: 428-436.
- Wang F, Nie J, Wang H, dkk. Characteristics of peripheral lymphocyte subset changes in COVID-19 pneumonia. *J Infeksi Dis*.
- Henry BM, de Oliveira MHS, Benoit S, Plebani M, Lippi G. Hematological, biochemical and immune biomarker abnormalities associated with severe illness and death in coronavirus disease 2019 (COVID-19): meta-analisis. *Clin Chem Lab Med (CCLM)*.
- Wang D, Hu B, Hu C, dkk. Clinical characteristics of 138 inpatients with pneumonia

- infected with the new coronavirus 2019 in Wuhan, China. JAMA. 2020; 323: 1061-1069.
- Zhao D, Yao F, Wang L, dkk. Comparative study of clinical features of Coronavirus 2019 pneumonia (COVID-19) with other pneumonia. Clin Infect Dis. 2020; 71 (15): 756-761.
- Zhang G, Hu C, Luo L, dkk. Clinical overview and short-term results of 221 patients with COVID-19 in Wuhan, China. J Clin Virol. 2020; 127: 104364.
- Liu K, Chen Y, Lin R, Han K. Clinical overview of COVID-19 in elderly patients: Comparisons with young and aged patients.
- Peng J, Qi D, Yuan G, dkk. Diagnostic value of peripheral hematology markers for coronavirus disease 2019 (COVID-19): studi multisenter, cross-sectional. J Clin Lab Anal. 2020; 34: e23475. Chen X, Yan L, Fei Y,.
- Zhang C. Laboratory abnormalities and risk factors associated with hospital deaths in patients with severe COVID-19. J Clin Lab Anal. 2020: e23467.